POTENTIAL ENHANCED
GREENHOUSE EFFECTS

STATUS AND OUTLOOK

DUANE G. LEVINE

PRESENTATION
TO
THE BOARD OF DIRECTORS
OF
EXXON CORPORATION

FEBRUARY 22, 1989

THE GREENHOUSE EFFECT IS REAL...HAS EXISTED THROUGHOUT MAN'S HISTORY...AND IN FACT...WITHOUT IT, CURRENT LIFE COULD NOT EXIST. TODAY'S CONCERNS ARE ABOUT AN ENHANCEMENT OF THIS EFFECT DUE TO HUMAN ACTIVITIES. So, I'LL REFER TO THESE CONCERNS COLLECTIVELY AS "POTENTIAL ENHANCED GREENHOUSE" OR PEG. IT HAS BEEN UNDER INTENSIVE SCIENTIFIC STUDY FOR OVER A DECADE BEFORE IT RECENTLY LEAPED TO THE FRONT PAGE.

IN SPITE OF THE RUSH BY SOME PARTICIPANTS IN THE GREENHOUSE DEBATE TO DECLARE THAT THE SCIENCE HAS DEMONSTRATED THE EXISTENCE OF PEG TODAY...I DO NOT BELIEVE SUCH IS THE CASE. ENHANCED GREENHOUSE IS STILL DEEPLY IMBEDDED IN SCIENTIFIC UNCERTAINTY, AND WE WILL REQUIRE SUBSTANTIAL ADDITIONAL INVESTIGATION TO DETERMINE THE DEGREE TO WHICH ITS EFFECTS MIGHT BE EXPERIENCED IN THE FUTURE.

POTENTIAL ENHANCED GREENHOUSE (PEG)

What's Known/Not Known About the SCIENCE

- Chemistry/Physics
- Climate Models
- Data
- Projections

Who Are the Principal PLAYERS

- Programs
- Plans
- Perceptions


What's NEXT

SO THIS REVIEW WILL BEGIN WITH WHAT IS KNOWN AND NOT KNOWN ABOUT THE SCIENCE...FIRST THE ESSENTIAL CHEMISTRY AND PHYSICS...AND THEN PROCEED WITH A DESCRIPTION OF CLIMATE MODELS WHICH ARE USED TO PREDICT FUTURE POTENTIAL

EFFECTS. I'LL COVER THE <u>HARD DATA</u> AND THEN SUMMARIZE <u>PROJECTED EFFECTS</u>...BASED ON COMPLEX CLIMATE MODELS...WHICH...INCI-DENTLY...HAVE YET TO BE VERIFIED.

However, policy initiatives are being advanced now and they could well out-pace scientific progress. I'll cover the current roster of principal players in this debate, outline their program and plans, and summarize some of their most visible perceptions. Finally, I'll try a glimpse at what's <u>next</u>.

GLOBAL ATMOSPHERIC CONCERNS




TO PUT PEG IN PERSPECTIVE, LET'S START WITH A GENERAL MODEL OF THE EARTH'S ATMOSPHERE AND 2 OF THE CURRENT MAJOR GLOBAL ATMOSPHERIC CONCERNS. ON THE LEFT, SOLAR RADIANT ENERGY IS REPRESENTED BY A BEAM RANGING FROM INFRARED (IR IN RED) TO ULTRAVIOLET (UV IN BLUE)...WITH THE BULK OF THE ENERGY IN THE VISIBLE RANGE SHOWN IN YELLOW. MOST OF THE

UV IS PICKED UP IN THE 03 LAYER IN THE STRATOSPHERE OR UPPER ATMOSPHERE (WHICH EXTENDS APPROXIMATELY 15-50 KM ABOVE THE EARTH'S SURFACE). THIS, OF COURSE, REVEALS THE CONCERN OVER DETERIORATION OF THIS LAYER BY MAN-MADE CHLOROFLUOROCARBONS (CFCs) WHICH CAN ATTACK 03 AND REDUCE ITS CONCENTRATION IN THE STRATOSPHERE. SUCH DETERIORATION WOULD ALLOW MORE UV TO PENETRATE THE ATMOSPHERE AND IMPACT LIFE ON THE EARTH'S SURFACE. PROTECTION OF THIS 03 LAYER BY REDUCING CFCs IS THE GOAL OF THE RECENT MONTREAL PROTOCOLS.

MOST OF THE IR PENETRATES THE STRATOSPHERE AND IS ABSORBED BY GREENHOUSE GASES IN THE LOWER ATMOSPHERE OR TROPOSPHERE. THIS, OF COURSE, HELPS WARM THE EARTH. THE BULK OF THE RADIANT ENERGY REACHING THE EARTH IS ABSORBED IN THE VISIBLE RANGE (SHOWN IN YELLOW)...DIRECTLY WARMING THE EARTH. THE RE-EMITTED ENERGY...AND THIS IS MOST IMPORTANT...IS CONCENTRATED IN THE IR RANGE AND THAT ADDS TO THE AMOUNT OF IR IN THE TROPOSPHERE.

OVERALL, THEN, THE UPPER ATMOSPHERE IS DOMINATED BY UV...AND 03 LAYER PROTECTION IS THE ISSUE...WHILE THE LOWER ATMOSPHERE IS DOMINATED BY IR AND GREENHOUSE-TYPE PROCESSES... WITH CLIMATE CHANGE AS THE ISSUE.

ATMOSPHERIC GREENHOUSE EFFECT




Troposphere

Earth Concentrates iR

WE CAN GO A LITTLE DEEPER ON GREENHOUSE TO HIGHLIGHT 2 KEY ASPECTS WHICH DETERMINE ITS OVERALL EFFECTIVENESS. AGAIN...THE BAND OF SOLAR RADIATION IS DEPICTED ON THE LEFT. AS MENTIONED, MOST OF THE INCIDENT VISIBLE SOLAR RADIATION (IN YELLOW) IS ABSORBED BY THE EARTH'S SURFACE AND RE-EMITTED ALMOST EXCLUSIVELY AS IR (IN RED). IF THIS RE-EMITTED IR ENCOUNTERS GREENHOUSE GAS-ES...LIKE CO2...WHOSE PROPERTIES FAVOR IR ABSORPTION...THEY ABSORB IT...LEVERAGING THE IR WARMING... INDICATED BY THE DASHED LINE. WHEN NO GREENHOUSE GASES ARE ENCOUNTERED, THE RE-EMITTED IR ESCAPES TO SPACE. THUS, THE EARTH SERVES TO CONCENTRATE IR IN THE TROPOSPHERE AND GREENHOUSE GASES CONVERT IT

ATMOSPHERIC GREENHOUSE EFFECT



Troposphere

- Earth Concentrates IR
- Water Vapor Increases IR Trapping Effect

TO HEAT. Now...THIS ADDITIONAL HEATING PROMOTES MORE WATER VAPORIZATION AND WATER VAPOR ITSELF IS A VERY EFFECTIVE GREENHOUSE GAS. IN FACT, IT IS ABOUT 3 TIMES MORE EFFECTIVE THAN CO2. It's really the water VAPOR THAT DOES MOST OF THE JOB. THIS LATTER POINT IS SIGNIFICANT BECAUSE ANY GREENHOUSE GAS...INCLUDING TRACE GREENHOUSE GASES LIKE CFCs, CH4, HALOGENS, N2O, AND O3 IN THE TROPOSPHERE...CAN TRIGGER THE ADDITIONAL WATER VAPOR LEVERAGING EFFECT.

THIS INSIGHT ALSO EXPOSES THE ENORMOUS DIFFERENCES BETWEEN 03 LAYER PROTECTION AND ENHANCED GREENHOUSE...

- THEY OCCUR IN DIFFERENT PARTS OF THE ATMOSPHERE.
- THEY INVOLVE DIFFERENT PHYSICS AND CHEMISTRY.
- THEY HAVE DIFFERENT EFFECTS ON LIFE.
 - WHEREAS 03 LAYER DETERIORATION IS CURRENTLY TRACED TO A SINGLE FAMILY OF MAN-MADE CHEMICALS...GREENHOUSE IN-CLUDES THE LIFE CYCLE GASES CO2 AND H20 WITH MANY NATURAL SOURCES AND SINKS.


FINALLY, I SHOULD ACKNOWLEDGE THAT THERE ARE SOME INTERACTIONS BETWEEN 03 LAYER PROTECTION AND GREENHOUSE IN THAT 03 AND CFCS ARE BOTH GREENHOUSE GASES AND 03 CAN REACT WITH CERTAIN OTHER GREENHOUSE TROPOSPHERIC GASES.

EXACTLY HOW THE GREENHOUSE GASES DETERMINE GLOBAL HEAT OR ENERGY BALANCE...AND HOW THE FLOW OF ENERGY IN AND OUT OF THE EARTH'S ATMOSPHERE DRIVES OUR CLIMATE IS DESCRIBED IN THE NEXT 2 FIGURES...FIRST, THE GLOBAL ENERGY BALANCE...

GREENHOUSE GASES & GLOBAL HEAT BALANCE

Radiation Balance Regulates Global Temperature

Cooling Depends On: Temperature Atmospheric Composition



VIEWED FROM SPACE, THE EARTH MUST BE IN ENERGY EQUILIBRIUM. THAT MEANS...HEAT FLOWING IN FROM SOLAR RADIATION (SHOWN IN YELLOW) MUST EQUAL HEAT RE-EMITTED BY THE EARTH (SHOWN IN RED). THIS PLOT SHOWS THE RADIANT ENERGY IN AND OUT AS A FUNCTION OF WAVE LENGTH...YELLOW SOLAR HEATING IN THE VISIBLE RANGE AND RED EARTH COOLING IN THE IR RANGE...ALTHOUGH THE SHAPES OF THE CURVES ARE DIFFERENT AND THE RADIATION IS AT QUITE DIFFERENT WAVELENGTHS, THE FACT IS THAT THE AREAS UNDER THE CURVES ARE EQUAL. THIS EQUALITY INDICATES THE BALANCE BETWEEN TOTAL RADIATIVE HEATING AND COOLING.


Now, incoming solar radiation...Is essentially a constant over time. However, radiant cooling depends on the Earth's temperature and atmospheric composition. The cooling curve is shown in more detail in the figure. Compared with the somewhat

GREENHOUSE GASES & GLOBAL HEAT BALANCE


Radiation Balance Regulates Global Temperature

Cooling Depends On:

Temperature Atmospheric Composition



Greenhouse Gases Selectively Trap IR Radiation



Existing Greenhouse Produces 60°F Surface Warming
 Without IR Greenhouse Earth Would Be Largely Frozen


IDEALIZED, SMOOTH COOLING CURVE SHOWN ABOVE...HERE, IN FINER DETAIL, WE SEE THE EFFECT OF GREENHOUSE GASES. THE CLEAR AREA BENEATH THE CURVE SHOWS THE ABSORPTION OF IR RADIANT ENERGY RE-EMITTED BY THE EARTH AT PARTICULAR WAVELENGTHS. QUITE OBVIOUS ARE THE STRONG ABSORPTION FEATURES PRODUCED BY CO2 AND O3. WATER VAPOR, THE MOST POWERFUL GREENHOUSE GAS, ABSORBS AT ALL WAVE LENGTHS. THE TOTAL GREENHOUSE EFFECT DEPENDS ON THE NATURE (MOLECULAR STRUCTURE) AND NUMBER (ATMOSPHERIC CONCENTRATION) OF ALL GREENHOUSE MOLECULES.

SINCE THE SOLAR HEATING INPUT REMAINS CON-STANT...WHILE THE GREENHOUSE GASES HAVE ABSORBED SOME OF THE IR COOLING, THE EARTH MUST INCREASE ITS TEMPERATURE TO MAINTAIN EQUILIBRIUM RADIANT HEAT OUTPUT. IN MODERN CLIMATE, THE EXISTING GREENHOUSE PRODUCES A 600F SURFACE WARMING. WITHOUT THIS EFFECT ... THE EARTH WOULD BE LARGELY FROZEN AND LIFE AS WE KNOW IT COULD NOT EXIST.

Now for the second figure...Describing how the flow of energy in the atmosphere drives climate.

ELEMENTS OF CLIMATE

Climate Response is Governed By Complex Interactions




ALTHOUGH OUR TOTAL ATMOSPHERE VIEWED FROM OUTSIDE SPACE MUST BE IN EQUILIBRIUM WITH RESPECT TO SOLAR HEATING AND EARTH COOLING, WIDE VARIATIONS IN HEATING CAN EXIST INSIDE OUR ENVIRONMENT. ONCE HEAT IS ABSORBED, IT SETS IN MOTION ALL THE COMPLEX PROCESSES WHICH SHAPE OUR CLIMATE. THE PURPOSE OF CLIMATE MODELS IS TO ATTEMPT TO SORT-OUT THESE EFFECTS. CLIMATE REFERS TO THE AVER-AGE TREND OF WEATHER, INCLUDING VARIABILITY. DYNAMIC EFFECTS OF WINDS AND CURRENTS CON-TROL GLOBAL CLIMATE BY TRANSPORTING HEAT OVER LARGE DISTANCES. IMPORTANT FEATURES OF AVERAGE CLIMATE INCLUDE EVAPORATION/ PRECIPITATION AND CLOUD FORMATION...ALL OF WHICH DISPLAY ENORMOUS VARIABILITY.

THE DIFFICULTY IN PREDICTING CLIMATE CHANGE...THEN...IS NOT ASSOCIATED WITH GREENHOUSE GAS IR RADIATIVE PROPERTIES...BUT RATHER WITH CAPABILITIES TO UNDERSTAND AND MODEL THE RESPONSE OF CLIMATE. FOR EXAMPLE, LET'S BRIEFLY OUTLINE THE EFFECTS OF A COMPOSITIONAL CHANGE...INCREASING CO2...ON CLIMATE.

ELEMENTS OF CLIMATE

• Climate Response is Governed By Complex Interactions




THE ADDED CO2 TRAPS SOME ADDITIONAL HEAT...WARMING THE ATMOSPHERE BY A SMALL AMOUNT. THIS TRIGGERS OTHER CHANGES. MODELS SHOW THAT A WARMER ATMOSPHERE BECOMES MORE MOIST...WITH MORE WATER VAPOR (A MORE POWERFUL GREENHOUSE GAS) CAUSING EVEN MORE WARMING. BUT DEPENDING ON CLIMATE RESPONSE...AN INCREASE IN MOISTURE MAY INCREASE CLOUD FORMATION...SHIELDING PARTS OF THE EARTH'S SURFACE FROM DIRECT SOLAR RADIATION, AND...PERHAPS...COMPENSATING FOR THE ADDITIONAL GREENHOUSE WARMING. OTHER EFFECTS NOT WELL UNDERSTOOD...LIKE OCEAN

CURRENTS...CAN ALSO AMPLIFY OR REDUCE WARMING.

In addition to atmospheric composition changes, we are also concerned today with climate changes forced by massive deforestation. This destruction adds to the CO2 Loading to the atmosphere and removes some CO2 sink capacity through photosynthesis.

FINALLY, OVER VERY LONG GEOLOGICAL TIME SCALES...CONTINENTAL DRIFT, SHIFTS IN THE EARTH'S ORBIT, AND OTHER SOLAR VARIATIONS...MAY CAUSE CHANGES IN SEA LEVEL AND SOLAR RADIATION WHICH CAN HAVE ENORMOUS EFFECTS ON CLIMATE. BUT THESE ARE USUALLY NOT PART OF CURRENT PEG PREDICTIONS.

HISTORICAL RECORD OF ATMOSPHERIC CO2 VARIATION



- Measurements Confirmed in Detail
- Increase 3 GTON Carbon Per Year
- Pre-Industrial CO₂ Level 270 ppm (28% Growth)


Now I'll cover the hard data we have on PEG. First, the famous Mauna Loa (Hawaii) recordings showing the increasing concentration of atmospheric CO2. Observations at The Scripps Institute began in 1958 as part of the International Geophysical Year. The data show a steady increase in CO2 with an obvious annual oscillation. The periodic variation occurs because CO2 is drawn down during the growing season and released in the fall and winter. Since 1958, CO2 has risen from 315 ppm to about 345 ppm, today.

THESE MEASUREMENTS HAVE BEEN CONFIRMED (± 1 PPM) IN GREAT DETAIL AT SITES FROM THE NORTH POLE TO ANTARCTICA. CO2 MIXES RAPIDLY IN THE ATMOSPHERE, SO RESULTS ON THE AVERAGE GROWTH OF CO2 ARE SIMILAR ACROSS THE GLOBE.

THE NET ACCUMULATION RATE OF CARBON IN THE ATMOSPHERE IS ABOUT 3 GIGATONS (BILLIONS OF METRIC TONS) PER YEAR CORRESPONDING TO THIS GROWTH.

FROM MEASUREMENTS OF AIR BUBBLES TRAPPED IN GLACIAL ICE CORES, IT IS ESTABLISHED THAT THE RECENT BUILDUP OF CO2 BEGAN WITH THE INDUSTRIAL ERA. IN THE MID 1800S THE CO2 LEVEL WAS ABOUT 270 PPM (* 10 PPM). CO2 GROWTH IS ABOUT 28% SINCE THEN.

HISTORICAL RECORD OF FOSSIL FUEL CO2 EMISSION



- Growth Resumed Following Reversal in Mid 70s

 Growth Rate Again at Historical Levels 4%/Yr
- If Growth Persists, CO₂ Doubling Occurs Sooner
 By Several Decades


THE NEXT DATA SHOW CO2 EMISSION FROM FOSSIL FUELS BETWEEN 1950 AND TODAY. THE UNITS ARE BILLIONS OF TONS (GTON) OF CARBON PER YEAR. THERE ARE TWO IMPORTANT POINTS FROM THIS FIGURE:

FIRST NOTE THAT MODERN EMISSION RATES ARE MUCH LARGER THAN THE RATE OF ACCUMULATION OF CO2 IN THE ATMOSPHERE. RECALL THAT ATMOSPHERIC CO2 IS GROWING BY ABOUT 3 GTON/YR. FOSSIL FUEL EMISSIONS TODAY ARE ABOUT 5 GTON, AND DEFORESTATION IS THOUGHT TO ADD ABOUT 1 GTON. EXACT VALUES FOR DEFORESTATION ARE CONTROVERSIAL. ESTIMATES RANGE FROM LESS THAN 1 TO OVER 2 GTON PER YEAR. NONETHELESS, ONLY ABOUT HALF THE EMITTED CO2 STAYS IN THE ATMOSPHERE. THE OTHER HALF IS THOUGHT TO BE ABSORBED INTO THE OCEANS. IT IS ALSO TRUE THAT PLANTS GROW BETTER IN AN ATMOSPHERE ENRICHED IN CO2. THAT HAS LED

SOME SCIENTISTS TO SUGGEST THAT CO2 IS BEING TAKEN UP AND CONVERTED TO BIOMASS. BUT THAT IS VERY DIFFICULT TO CONFIRM IN STUDIES OF NATURAL ECOLOGICAL SYSTEMS SO FAR.

THE SECOND POINT FROM THE FIGURE CONCERNS THE GROWTH IN CO2 EMISSION. PRIOR TO THE ARAB OIL EMBARGO, EMISSIONS GREW STEADILY AT ABOUT 4.4% PER YEAR. WHEN THE GREENHOUSE ISSUE WAS FIRST IDENTIFIED IN THE MID-1970s, THAT RATE WAS EXTRAPOLATED TO PROJECT ATMO-SPHERIC CO2 LEVELS WOULD DOUBLE EARLY IN THE NEXT CENTURY, SAY 2025. HOWEVER, THE RE-DUCED USE OF FOSSIL FUELS SINCE THEN, CAUSED FORECASTS OF DOUBLING TO MOVE OUT BY SEVERAL DECADES, SAY TILL 2075-2100. THE FIGURE SHOWS THAT THE GROWTH RATE HAS ONCE AGAIN RECOVERED TO PAST HISTORICAL LEVELS. HIGHER GROWTH PERSISTS, THE DOUBLING TIME WILL AGAIN MOVE CLOSER BY SEVERAL DECADES.

HISTORICAL RECORD OF GLOBAL TEMPERATURE CHANGE



Noteworthy Differences With Greenhouse Predictions

- Recent Warming Reverses Cooling From 1940-1970s
 - 1980s Warmest Decade on Record
 - Persistent Trend Could Signal Greenhouse Warming
- US 1988 Summer Heat and Drought, A Critical Media Event
 - Not a Predicted Consequence of Greenhouse Warming
 - Due to a Natural Weather Fluctuation
 - Cited as an Example of Future Trends

THE FINAL DATA SET COVERS PAST VARIATION IN GLOBAL AVERAGE TEMPERATURE. OF COURSE, GLOBAL AVERAGE TEMPERATURE IS A STATISTICAL CONCEPT. IT CANNOT BE DIRECTLY MEASURED. THERE ARE SERIOUS ISSUES CONCERNING COMPLETENESS, ACCURACY, AND INTERPRETATION OF THIS HISTORICAL DATA. HOWEVER, THE PROBLEM IS RECEIVING A GREAT DEAL OF ATTENTION, AND MOST STUDIES SHOW RESULTS SIMILAR TO THESE.

THE DATA SHOW A RELATIVELY LARGE SCATTER OF A FEW TENTHS OF A DEGREE FROM YEAR TO YEAR. THIS "NOISE" OCCURS FROM NATURAL FLUCTUATIONS THAT ARE NOT COMPLETELY UNDERSTOOD. IT IS KNOWN THAT EVENTS LIKE VOLCANIC ERUPTIONS AND CHANGES IN OCEANIC UPWELLING (SUCH AS EL NINO), CAUSE PART OF THE VARIABILITY. THE DASHED TREND LINE ILLUSTRATES THE GENERAL BEHAVIOR. THE RECORD SHOWS AN APPARENT RISE OF ABOUT 1/20 C OVER THE PAST 100 YEARS.

However, the warming does not agree with models based on CO2 variations. In particular, enhanced Greenhouse models predict a smoothly accelerating increase of temperature with time (shown in red). The data are quite different. Most noticeable is a cooling trend between the 1930s and late 1970s when the model predicts warming.


DATA ON TEMPERATURE VARIATION IN THE 1980s ARE NOW BECOMING AVAILABLE. THEY SHOW A REVERSAL OF THE RECENT COOLING TREND. IN FACT, THE 3 WARMEST YEARS ON RECORD OCCURRED IN THE 1980s. IF THIS TREND PERSISTS IT

COULD SIGNAL THAT ENHANCED GREENHOUSE WARMING IS FINALLY BECOMING DETECTABLE.

THERE IS NO DOUBT THAT THE 1988 HEAT AND DROUGHT WERE A CRITICAL EVENT IN THE GREEN-HOUSE ISSUE, BECAUSE THEY STIRRED PUBLIC ATTENTION, AND BROUGHT HOME POTENTIAL CONSEQUENCES OF CLIMATE WARMING. HOWEVER, 1988 WAS NOT A PREDICTED CONSEQUENCE OF ENHANCED GREENHOUSE MODELS. MOST METEOROLOGISTS INTERPRET THE SUMMER AS AN INFREQUENT, BUT NOT UNEXPECTED FLUCTUATION IN WEATHER. GREENHOUSE SCIENTISTS BY AND LARGE HAVE NOT CLAIMED THE US HEAT AND DROUGHT WERE CAUSED BY ENHANCED GREENHOUSE, BUT THEY HAVE CITED IT AS AN EXAMPLE OF WHAT PEG MIGHT BRING.

So far we have discussed the historical record, and shown the important data. However, PEG impacts occur in the future. So now we turn to projections.


PROJECTED SOURCES OF ENERGY AND CO2 EMISSION



- Future Emission Sensitive to Forecasts of Energy Demand, Source
 - Projections Differ Significantly Beyond Near Term

FORECASTS OF FUTURE LEVELS OF ATMOSPHERIC CO2 BEGIN BY PROJECTING ENERGY NEEDS AND SOURCES WELL INTO THE NEXT CENTURY. THE FIGURE SHOWS THREE FORECASTS OF ENERGY DEMAND TAKEN FROM A DOE STUDY, LABELED CASE A, B, AND C. THE UNIT OF ENERGY IS 1018 JOULES, WHICH IS ABOUT THE SAME AS QUADS, OR QUADRILLIONS OF BTUS. CASE B IS CLOSE TO RECENT EXXON PROJECTIONS.

PROJECTED SOURCES OF ENERGY AND CO2 EMISSION




Future Emission Sensitive to Forecasts of Energy Demand, Source
 Projections Differ Significantly Beyond Near Term

IT IS WELL KNOWN THAT FOSSIL FUELS EMPLOYED TO SATISFY THE DEMANDS, DIFFER IN THE AMOUNT OF CO2 RELEASED TO PRODUCE A BTU OF ENERGY. THIS PART OF THE FIGURE COMPARES FOSSIL FUELS ON A RELATIVE SCALE WHERE EMISSIONS PER BTU FROM CONVENTIONAL OIL ARE TAKEN AS ONE. THEN, GAS IS 0.7 AND COAL IS 1.25. WE ALSO SHOW THE INDEX VALUES FOR SYNTHETIC LIQUIDS. SYNTHETIC LIQUIDS FROM COAL AND SHALE RESULT IN GREATER CO2 EMISSIONS. THE EXTRA CO2 IS GENERATED IN THE MANUFACTURING PROCESS. ESPECIALLY FOR SYNTHETICS FROM

Synthetic Liquids

COAL AND SHALE THE CO2 FACTOR IS SENSITIVE TO THE RAW MATERIAL AND PROCESS ROUTE. LIQUIDS DERIVED DIRECTLY FROM GAS HAVE SIGNIFICANTLY LOWER CO2 INDICES.

PROJECTED SOURCES OF ENERGY AND CO2 EMISSION




Future Emission Sensitive to Forecasts of Energy Demand, Source
 Projections Differ Significantly Beyond Near Term

FOR EACH ENERGY DEMAND CASE...(A, B, & C)...
THIS CO2 GENERATION INDEX CAN BE USED TO
COMPUTE CO2 EMISSIONS AS SHOWN IN THE FINAL
FIGURE. CASE A WITH THE GREATEST ENERGY
DEMAND ALSO PRODUCES MUCH HIGHER CO2 EMISSIONS, SINCE IT RELIES ON HIGH CO2 PRODUCING
FUELS FROM SYNTHETICS. OVERALL FORECASTS
DIFFER WIDELY BEYOND THE NEAR TERM. EXTENDED OUT TO 2100 THEY VARY BY FACTORS OF 20 IN
CO2 EMISSION LEVELS.

PROJECTED GROWTH OF ATMOSPHERIC CO2

Combine Forecasts: Economic Growth, Energy Use, CO₂ Growth



- Atmospheric CO₂ Doubles (600 ppm)
 - Case A: 2030
 - Case B: 2060
 - Case C: 2100

BY COMBINING THESE THREE TYPES OF PROJECTIONS FOR THE FUTURE...

- TOTAL ENERGY CONSUMPTION
- CO2 EMISSION FACTORS FOR VARIOUS FOSSIL FUELS

AND

 DISTRIBUTION OF FOSSIL FUEL ENERGY CON-SUMPTION BY RESOURCE...

IT IS POSSIBLE TO FORECAST FUTURE LEVELS OF ATMOSPHERIC CO2. THE PROJECTIONS SHOW DOE FORECASTS FOR THE 3 CASES THROUGH THE YEAR 2080.

IN THESE FORECASTS THE CO2 DOUBLING TIME IS A CONVENIENT BENCHMARK TO MEASURE CO2 BUILD-UP. (IT IS THE LEVEL CONSIDERED IN MOST CLIMATE SIMULATIONS OF THE GREENHOUSE EFFECT.) However, There is nothing magical about doubling itself. Taking the doubling Level to be 600-700 ppm, the various cases LEAD to a doubling between 2030 and 2100 as INDICATED.

POTENTIAL CLIMATE IMPACT FROM CO2: NEXT 100 YEARS

"CHANGING CLIMATE", NATIONAL RESEARCH COUNCIL 1983

- Temperature
 - Global Mean Temperature Increase ... 1.5 4.5°C
 - Greater Warming in Polar Regions (2-3x)
- Sea Level/Sea Ice
 - Coverage and Thickness of Sea Ice/Glaciation Will Decrease
 - Sea Level Rise (Meltwater + Thermal Expansion) ... 70 cm
- Natural Ecosystems and Agriculture
 - Regional Climate Change: Temperature, Hydrology
 - Enhanced Productivity From Increased CO₂
 - Global Net Effect Uncertain

THESE CO2 PROJECTIONS ARE USED IN CURRENT CLIMATE MODELS TO PREDICT IMPORTANT CHANGES OVER THE NEXT 100 YEARS. THIS SET OF RESULTS IS TAKEN FROM THE NATIONAL RESEARCH COUNCIL (NRC) REPORT "CHANGING CLIMATE".

Consensus predictions call for warming between 1.5-4.5 oc for doubled CO2 with greater warming at the poles. Note that these numbers reflect the range produced by available models. No one knows how to evaluate the absolute uncertainty in the numbers.

THE EXTENT AND THICKNESS OF GLACIERS ARE PREDICTED TO DECREASE, LEADING TO SEA LEVEL RISE. THE NRC REPORT CHOSE A MOST LIKELY VALUE OF 70 CM SEA LEVEL RISE. OTHER PREDICTIONS SUGGEST A BROADER RANGE FROM 30-200 CM. THE RISE OCCURS BOTH FROM A LARGER AMOUNT OF WATER IN THE OCEANS, AND FROM THERMAL EXPANSION.

FINALLY, CLIMATE CHANGE AND HIGHER LEVELS OF ATMOSPHERIC CO2 AFFECT AGRICULTURE AND ECOSYSTEMS. THERE ARE TWO ASPECTS. FIRST, THE DIRECT EFFECT OF CLIMATE CHANGE ALTERS THE LENGTH OF THE GROWING SEASON AND THE AVAILABILITY OF WATER. SECOND, NEARLY ALL PLANTS GROW MORE RAPIDLY, AND USE LESS WATER, IN A HIGH CO2 ENVIRONMENT. THE SECOND EFFECT CAN BE QUITE POSITIVE FOR MANAGED AGRICULTURE.

MODELS CANNOT YET PREDICT REGIONAL CLIMATE CHANGE WITH MUCH ACCURACY. I AM SURE YOU HAVE ALL HEARD THAT THE US MID-WEST MAY BECOME A DUST BOWL FROM ENHANCED GREENHOUSE, WHILE RUSSIA MAY BECOME MORE FRUITFUL. THAT IS A PROJECTION OF SOME MODELS, BUT OTHERS SHOW THE OPPOSITE.

THE CLIMATE MODEL IMPACTS OF PEG ARE NOT ALL NEGATIVE. THEY AFFECT DIFFERENT PARTS OF THE GLOBE UNEQUALLY. THE MODELS PREDICT VARIOUS WINNERS AND LOSERS.

THIS COMPLETES THE GREENHOUSE SCIENCE DISCUSSION.

SUMMARIZING:

DATA CONFIRM THAT

- GREENHOUSE GASES ARE INCREASING IN THE ATMOSPHERE.
- Fossil fuels contribute most of the CO2...But deforestation is also significant.

AND

• HISTORICAL TEMPERATURES SHOW ONLY SLIGHT WARMING...NOT ENOUGH TO CONFIRM ENHANCED GREENHOUSE.

THAT'S ALL THE DATA.

PROJECTIONS SUGGEST

 Significant climate change with a variety of regional impacts.

AND

 SEA LEVEL RISE WITH GENERALLY NEGATIVE CONSEQUENCES.

FINALLY, THE SIZE AND TIMING OF IMPACTS ARE UNCERTAIN.

POTENTIAL ENHANCED GREENHOUSE

PLAYERS...

0 US

CONGRESS/PRESIDENT EPA/DOE/STATE NOAA/NASA/NSF

AND THE MEDIA

0 <u>III</u>

GENERAL ASSEMBLY - "WORLD COMMISSION ON ENVIRONMENT AND DEVELOPMENT" UNEP/NMO

- O CANADA
 PRIME MINISTER
 FEDERAL ENVIRONMENTAL MINISTRY
- 6 INTERNATIONAL COUNCIL OF SCIENTIFIC UNIONS
- EEC
- OECD
- ENVIRONMENTAL GROUPS

THE KEY PLAYERS ON PEG ARE IN THE US AND UN. IN THE US, CONGRESS IS VERY INTERESTED. SENATORS LIKE WIRTH, MITCHELL, BAUCUS, AND GORE HAVE VERY HIGH PROFILES. PRESIDENT BUSH HAS COMMITTED TO CONVENE A GLOBAL GREENHOUSE CONFERENCE. WITHIN THE ADMINISTRATION, THE PLAYERS ARE EPA (WHICH HAS A PARTICULARLY CRITICAL ROLE THROUGH ITS INTERNATIONAL ACTIVITIES), DOE (WHICH SPONSORED AN EXTENSIVE REVIEW OF THE SUBJECT COMPLETED IN 1985), AND THE STATE DEPARTMENT (WHOSE INTERESTS ARE DIRECTED TOWARDS INTERNATIONAL TREATIES AND CONVENTIONS). ADDITION, TECHNICAL AGENCIES INCLUDING NOAA, NASA, AND NSF HAVE SIGNIFICANT ROLES...AND, OF COURSE, THE MEDIA IS VERY ACTIVE.

ALL OF THESE EFFORTS ARE ESTABLISHING LINKS WITH OTHER NATIONAL AND INTERNATIONAL INTER-ESTS...ESPECIALLY IN THE UN WHICH HAS DEVOT-ED A GREAT DEAL OF EFFORT TO PEG. THE GENERAL ASSEMBLY ESTABLISHED A WORLD-WIDE COMMISSION HEADED BY MME. GRO BRUNDTLAND (PRIME MINISTER OF NORWAY) TO EVALUATE PROBLEMS OF ENVIRONMENTAL PROTECTION AND ECONOMIC DEVELOPMENT. THE COMMISSION'S REPORT... "OUR COMMON FUTURE"... CONTAINS A HEAVY DOSE OF GREENHOUSE CONCERNS...ON CLIMATE, AGRICULTURE, ENERGY...AND LISTS IT AS THE #1 ENVIRONMENTAL PROBLEM WE FACE. THE GENERAL ASSEMBLY HAS DIRECTED ALL OF ITS AGENCIES TO DEVELOP SPECIFIC PLANS TO DEAL WITH PEG.

THE UNITED NATIONS ENVIRONMENTAL PROGRAM (UNEP) IS ONE OF THESE AGENCIES AND IT HAS

MADE PEG ITS #1 PRIORITY. UNEP...LIKE EPA HAS A CRITICAL ROLE...HAVING BEEN DIRECTED TO DEVELOP DETAILED PLANS WITH THE WORLD METEOROLOGICAL ORGANIZATION (WMO) FOR ACTION PROGRAMS AND POLICY PROPOSALS TO BE RECOMMENDED TO THE GENERAL ASSEMBLY.

THERE ARE NUMEROUS OTHER PLAYERS. THE CANADIANS, INCLUDING THEIR PRIME MINISTER AND FEDERAL ENVIRONMENTAL MINISTRY, HAVE BEEN...PERHAPS...THE MOST OUTSPOKEN GOVERNMENT IN RESPONDING TO THE NEED FOR GREENHOUSE LIMITATIONS. THEY SPONSORED A WORLD CONFERENCE ON PEG DURING THIS PAST JUNE...WHICH CALLED FOR 20% REDUCTION IN CO2 EMISSIONS BY 2010 WITH AN OVERALL GOAL OF 50% REDUCTION...AND A TAX ON FOSSIL FUELS TO FUND ALTERNATE ENERGY DEVELOPMENT.

THE INTERNATIONAL COUNCIL OF SCIENTIFIC UNIONS (ICSU) IS PROVIDING A FORUM FOR WORLD-WIDE SCIENTIFIC EFFORTS UNDER THE BANNER OF THE "INTERNATIONAL BIOPHYSICAL YEAR". THE EUROPEAN ECONOMIC COMMUNITY (EEC) IS PURSUING ITS POSITION AND DEVELOPED COUNTRIES IN THE OECD HAVE ESTABLISHED ACTIVITIES TO PARTICIPATE IN THE EXPECTED DEBATE.

OF COURSE, NUMEROUS ENVIRONMENTAL GROUPS...INCLUDING HIGH PROFILE ORGANIZATIONS LIKE THE GREENS, WORLD RESOURCES INSTITUTE, AND NATIONAL RESOURCES DEFENSE COUNCIL...HAVE NOW INVESTED A LOT OF TIME AND EFFORT IN EXPERTISE RELATED TO ENERGY AND GREENHOUSE. HAVING INVESTED SIGNIFICANT

EFFORT, THEY NEED...AND INTEND...TO HAVE SIGNIFICANT IMPACT.

SCHEDULED ACTIVITIES - PEG

US		
<u>03</u>		
•	INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE (WITH WMO/ICSU)	1988
•	EPA REPORTS ON - POTENTIAL CONSEQUENCES FOR US/MORLD - MITIGATION/STABILIZATION APPROACHES	1988
•	DOE - ASSESSMENT OF RED ON ALTERNATE EMERGY - INVENTORY OF GREENHOUSE GASES - ANALYSIS OF PRIVATE SECTOR OPTIONS - POLICY OPTIONS TO LINIT EMISSIONS	1989
UN B	US/INTERNATIONAL SCIENCE ASSESSMENT	1990
•	COORDINATE WORLDWIDE SCIENTIFIC ASSESSMENT WHO/ICSU	1990
•	REGIONAL IMPACT ASSESSMENTS	
	POLICY COTTONS TO LEVER CO.	1990
•	POLICY OPTIONS TO LIMIT CLIMATE CHANGE	1990
•	INTERNATIONAL CONVENTION TO LIMIT CITMATE CHANCES	HTD 10011

THE INTENSITY OF THESE EFFORTS IS REFLECTED IN THE SLATE OF ACTIVITIES CURRENTLY SCHED-ULED. FOR EXAMPLE, IN '88 THE US, THROUGH NOAA AND NASA, SET UP AN INTERGOVERNMENTAL CLIMATE PANEL WITH WMO AND ICSU TO PROMOTE LIAISON BETWEEN GOVERNMENTS...AND EPA WAS TO REPORT TO CONGRESS ON BOTH POTENTIAL CONSEQUENCES OF PEG FOR THE US/WORLD AND... MITIGATION/STABILIZATION APPROACHES TO LIMIT CLIMATE CHANGE. THIS REPORT IS NOW DUE IN 1989. DOE WILL BE VERY BUSY. IN 1989, IT

IS DUE TO DEVELOP REPORTS ON VARIOUS ASPECTS OF PEG INCLUDING:

- Assessing alternate energy R&D
- CATALOGUING GREENHOUSE GASES
- ANALYZING ENERGY OPTIONS FOR THE PRI-VATE SECTOR

AND

 EVALUATING POLICY OPTIONS FOR LIMITING CO2 EMISSIONS

FINALLY, NSF IS PLANNING A US/INTERNATIONAL SCIENCE ASSESSMENT BY 1990.

THE UN HAS RECENTLY COMMISSIONED A WORLDWIDE SCIENTIFIC ASSESSMENT DESIGNATING WMO TO WORK CLOSELY WITH ICSU. BY THE WAY, NSF WILL ALSO COORDINATE CLOSELY WITH THIS EFFORT IN DEVELOPING ITS VIEWS. REGIONAL IMPACT STUDIES ARE BEING SET UP USING CLI-MATE MODELS TO PROJECT THE VARIOUS WINNERS AND LOSERS AS RAINFALL, WIND, STORMS, AND SEA LEVEL PATTERNS CHANGE. THIS COULD BE ESPECIALLY CONTENTIOUS BECAUSE OF THE WIDE ROOM FOR INTERPRETATION IN APPLYING THE MODELS AND THE ENORMOUS POTENTIAL POLITICAL CONSEQUENCES. CONCURRENTLY, UNEP WILL DEVELOP ITS VIEW OF POLICY OPTIONS TO LIMIT CLIMATE CHANGE. ALL OF THESE ACTIVITIES ARE PLANNED FOR COMPLETION IN 1990. UNEP HAS BEEN URGED TO AIM FOR AN INTERNATIONAL CONVENTION ON PEG BY ABOUT 1995. HOWEVER, THIS TARGET DATE SEEMS TO BE ADVANCING TOWARDS 1992.

GIVEN, THE ...

- COMPLEXITY OF THE SCIENCE
- ENORMOUS POTENTIAL GLOBAL IMPACTS
- DIVERSITY OF THE PLAYERS

AND

• INTENSITY OF THEIR ACTIVITIES...

WHERE IS ALL THIS HEADED? I BELIEVE THERE IS A PATTERN...AND IT'S ROOTED IN THE EVOLUTION OF THE JUST-COMPLETED MONTREAL PROTOCOLS TO PROTECT THE STRATOSPHERIC 03 LAYER BY LIMITING MAN-MADE CFCs.

STRATOSPHERIC OZONE/PEG ANALOGY

OZONE LAYER

ENHANCED GREENHOUSE

'74	· 🗸	ATMOSPHERIC CHEM/PHYS		
	4	GROWTH IN [CFC'S] / [CO2] & [TRACE GASES]	1	'75
	1	INDUSTRIAL SOURCES	1	
		MODELS: END EFFECT PROJECTIONS CONCEPT OF 'DELAY'	1	
	•	ENVIRONMENTAL CAUSE	4	
	4	INTERNATIONAL OWNERSHIP	1	
	y	US/UN AXIS	1	
	4	CRITICAL EVENT CALL FOR ACTION	1	
	VIENNA	CONVENTION	7	
'87		AL PROTOCOLS	?	
		+	γ	

ABOUT 20 YEARS AGO, THE ATMOSPHERIC CHEMISTRY AND PHYSICS BEGAN WITH CONCERNS OVER THE EFFECT OF SUPERSONIC TRANSPORTS ON THE 03 LAYER. AS THESE FEARS ABATED...IN 1974... SOME SCIENTISTS' LABORATORY TESTS INDICATED THAT THE VERY STABLE CLASS OF MAN-MADE CHEMICALS (CFCs) BREAK DOWN UNDER THE KIND

OF INTENSE UV RADIATION THAT IS PRESENT IN THE STRATOSPHERE...WITH THE RESULTING CL AND BR ATOMS ATTACKING AND DESTROYING 03.

SUBSEQUENTLY, IT WAS ESTABLISHED THAT THE CFC CONCENTRATIONS WERE INCREASING IN THE STRATOSPHERE. THESE CFCs, OF COURSE, WERE EASILY TRACED TO MAN-MADE SOURCES AND INDUSTRIAL APPLICATIONS.

EXTENSIVE MODELING EXERCISES PREDICTED...

- THE LONG TERM RATE OF 03 LAYER DETERI-ORATION
- THE RESULTING INCREASED UV PENETRATION OF THE EARTH'S ATMOSPHERE

AND

• THE SERIOUS POTENTIAL REPERCUSSIONS LIKE INCREASED HUMAN CANCER RATES AND PLANT DAMAGE

Now these predicted effects were well into the future. So a crucial step was the introduction of "a delay concept" based on the unusual chemical stability of CFCs in the lower atmosphere and the very long transport times for CFCs to reach the upper stratosphere. The reasoning was...there was an <u>already</u> committed 03 layer deterioration based on CFCs <u>already</u> "in the pipeline". This gave rise to an environmental cause which was quickly adopted as an international issue. The players were basically similar to those organizing around the current

GREENHOUSE ISSUE...PRIMARILY IN THE US AND UN. WHEN THE US BECAME ACTIVELY INVOLVED...INITIATING COOPERATION WITH THE UN
TO LIMIT WORLDWIDE CFC PRODUCTION AND
SALES...ALL OF THE ELEMENTS WERE IN PLACE.

BUT WITH ALL OF THIS, PROGRESS BEGAN TO LANGUISH AND THE EFFORT MIGHT WELL HAVE FOUNDERED, EXCEPT FOR THE DISCOVERY OF THE SO-CALLED "03 LAYER HOLE" OVER ANTARCTICA. THIS WAS A MOST CRITICAL EVENT - ALTHOUGH ITS EXACT RELEVANCE TO CFC RELATED 03 LAYER DETERIORATION REMAINS UNEXPLAINED. IT RE-ENERGIZED THE EFFORT AND DIRECTLY LED TO A "CALL FOR ACTION" AND THE VIENNA CONVEN-TION. SHORTLY THEREAFTER...IN 1987...THE MONTREAL PROTOCOLS TO LIMIT CFCs WITH A PHASED 50% REDUCTION BY THE TURN OF THE CENTURY WERE APPROVED AND ARE EXPECTED TO BE ADOPTED. I SHOULD ADD THAT IN 1988...JUST WITHIN THE PAST 6 MONTHS OR SO...THERE IS FOR THE FIRST TIME CONVINCING SCIENTIFIC EVIDENCE THAT 03 LAYER DETERIORATION HAS BEEN DETECTED. SO THAT AFTER THE FACT...SOME ACTION SEEMS JUSTIFIED.

Now, I hope from the material covered on Greenhouse so far...It is clear that we have advanced through similar stages. A critical event occurred with the "long hot summer of '88". Although most responsible scientists believe this was due to natural fluctuations in weather patterns...It has drawn much attention to the potential problems and we're starting to hear the inevitable call for action. Exactly what happens now is not

CLEAR...BUT THIS CRITICAL EVENT HAS ENERGIZED THE GREENHOUSE EFFORT AND RAISED PUBLIC CONCERN OVER PEG.

SOME KEY PERCEPTIONS/MISCONCEPTIONS...

- "PEG" IS PART OF/CLOSELY LINKED TO "03 LAYER PROTECTION"
- ENOUGH RESEARCH HAS BEEN DONE
 - EXISTENCE OF PEG IS NOW ESTABLISHED
 - ADVANCING THE SCIENCE WILL TAKE TOO LONG
- '88 SUPPLER WAS DUE TO PEG
- · CAN'T TOLERATE DELAY
- . NEED POLICY DEVELOPMENT NOW
- PROJECTED SHIFTS IN WEATHER WILL FEATURE BIG WINNERS AND LOSERS WITH MAJOR POLITICAL IMPLICATIONS
- DEVELOPED COUNTRIES WILL SHIFT RESOURCES TO DEVELOPING COUNTRIES
- O NUCLEAR AND/OR RENEMABLE ENERGY RESOURCES CAN SOLVE THE PROBLEM

OVERALL ...

• RISK FROM PEG CLIMATE CHANGES IS NOT ACCEPTABLE...
MUST DEVELOP (IMPLEMENT) CONTROL POLICIES NOW

In facing this reality, there are some key perceptions and misconceptions we'll have to deal with. The first misconception is that PEG and 03 layer are somehow the same problem... or at least closely linked. This is particularly dangerous because the two problems are, in fact, very different and effective solutions are bound to be very different... banning a single class of man-made CFC chemicals is very difficulti... but pales by comparison to the difficulties of applying similar approaches to life cycle gases like CO2 and water.

THE SECOND MISCONCEPTION IS THAT ENOUGH RESEARCH ON THE BASIC PROBLEM HAS BEEN DONE. FAILURE TO UNDERSTAND THE NEED FOR SUBSTANTIAL ADVANCES IN THE SCIENCE TO REDUCE THE UNCERTAINTY AND EXTREME VARIABILITY IN THE PROJECTIONS CAN LEAD TO PREMATURE LIMITATIONS ON FOSSIL FUELS.

Using the '88 summer weather patterns as part of this argument...the third misconception...is simply incorrect.

ARGUMENTS THAT WE CAN'T TOLERATE DELAY AND MUST ACT NOW CAN LEAD TO IRREVERSIBLE AND COSTLY DRACONIAN STEPS.

PROJECTING PEG INDUCED CLIMATIC REGIONAL WINNERS AND LOSERS...AND COMPETITION BETWEEN DEVELOPED AND DEVELOPING COUNTRIES IN RATIONING ENERGY TO LIMIT PEG...COULD EASILY DOMINATE ANY DEBATE AND MAKE RATIONAL CO-OPERATION LESS LIKELY.

THERE HAVE BEEN DRAMATIC SHIFTS IN ATTITUDES TOWARDS NUCLEAR ENERGY BY ENVIRONMENTAL GROUPS BECAUSE OF THEIR CONCERNS OVER PEG. FURTHERMORE, RENEWABLE ENERGY ADVOCATES HAVE TRADITIONALLY OVERSTATED CAPABILITIES. THESE BOTH TEND TO ENCOURAGE A PRECIPITOUS SHIFT TO ALTERNATE ENERGY AND UNDERSTATE THE CONSIDERABLE DIFFICULTIES WHICH MUST BE OVERCOME.

WHAT'S NEXT

Expect:

- Continued Pressure to...
 OVERSTATE SCIENTIFIC UNDERSTANDING
- Prominent Media Role to...
 INCREASE PUBLIC CONCERNS



More Initiatives to...
INCREASE PACE OF INTERNATIONAL NEGOTIATIONS

Rational Responses Require:

- Efforta to...
 EXTEND THE SCIENCE
- Emphasis on...
 COSTS/POLITICAL REALITIES



AS FOR "WHAT'S NEXT"...WE CAN EXPECT CONTIN-UED PRESSURE TO OVERSTATE CURRENT SCIENTIFIC UNDERSTANDING. THE MEDIA ROLE...ALREADY PROMINENT...IS LIKELY TO INCREASE PUBLIC AWARENESS AND CONCERN, AND THERE WILL BE CONTINUING INITIATIVES TO EXTEND INTERNA-TIONAL NEGOTIATIONS. AS THE DEGREE OF THESE EFFORTS EXCEED UNDERSTANDING (OR ABILITY TO RESPOND CONSTRUCTIVELY)...THERE IS A TENDEN-CY TOWARDS A "CRISIS MENTALITY." MEANWHILE, MORE RATIONAL RESPONSES WILL REQUIRE EFFORTS TO EXTEND THE SCIENCE AND INCREASE EMPHASIS ON COSTS AND POLITICAL REALITIES TO FRAME "ADAPTIVE" MEASURES WHICH ARE DOABLE AND MOVE TOWARDS CONSTRUCTIVE OPTIONS.

EXXON'S POSITION

IMPROVE UNDERSTANDING

Extend the Science Include the Costs/Economics Face the Socio-Political Realities

• STRESS ENVIRONMENTALLY SOUND ADAPTIVE EFFORTS

Support Conservation
Restrict CFCs
Improve Global Re/De Forestation

TO BE A RESPONSIBLE PARTICIPANT AND PART OF THE SOLUTION TO PEG, EXXON'S POSITION SHOULD RECOGNIZE AND SUPPORT 2 BASIC SOCIETAL NEEDS. FIRST...TO IMPROVE UNDERSTANDING OF THE PROBLEM...NOT JUST THE SCIENCE...BUT THE COSTS AND ECONOMICS TEMPERED BY THE SOCIO-POLITICAL REALITIES. THAT'S GOING TO TAKE YEARS (PROBABLY DECADES). BUT THERE ARE MEASURES ALREADY UNDERWAY THAT WILL IMPROVE OUR ENVIRONMENT IN VARIOUS WAYS...AND IN ADDITION REDUCE THE GROWTH IN GREENHOUSE GASES. THAT'S THE SECOND NEED INCLUDING THINGS LIKE ENERGY CONSERVATION, RESTRICTION OF CFC EMISSIONS, AND EFFORTS TO INCREASE THE GLOBAL RATIO OF RE/DE FORESTATION. OF COURSE, WE'LL NEED TO DEVELOP OTHER RESPONSE OPTIONS...IMPLEMENTING MEASURES WHEN THEY ARE COST EFFECTIVE IN THE NEAR TERM AND PURSUING NEW TECHNOLOGIES FOR THE FUTURE.